

Bufomicrus squamosus Sharp

Bufomicrus squamosus Sharp, 1891:145.

Distribution. British Honduras: "British Honduras (Blancaneaux)." *Guatemala:* Telemán, La Tinta, Chacoj, Tamahu, San Juan, Panima, all in the Polochic Valley, Vera Paz. Additional localities: El Estor, Estado de Izabal (on north side of Lago de Izabal), I-15-46, ELS, (ELS), 3 examples.

Bufomicrus globipennis Sharp

Bufomicrus globipennis Sharp, 1891:146.

Distribution. Guatemala: Yzabal (Salle). Known only from the two examples cited in the original description. It is probable that the locality "Yzabal" and Izabal are the same.

ACKNOWLEDGEMENTS

I am indeed grateful to Dr. Fred Truxal and the Los Angeles County Museum for making otherwise unobtainable literature available, and to Sir G. A. K. Marshall of the Commonwealth Institute of Entomology, British Museum (Natural History), for efforts on his part in aiding to properly place some of this and other material, and for pertinent information regarding the types.

LITERATURE CITED

- BUCHANAN, L. L. 1939. The species of *Pantomorus* of America north of Mexico. Misc. Publ. United States Dept. Agric. no. 341:1-39. Illus.
- SHARP, D. 1891. *Biologia Centrali-Americana*, Insecta, Coleoptera, Curculionidae: Apioninae, Thecesterninae, Otiiorhynchinae, vol. 4, pt. 3:81-168. Illus.
- SLEEPER, E. L. 1953. New genera and species of Curculionidae with a new species of Anthribidae (Coleoptera). *Ohio Journal. Sci.*, 53(2):113-120. Illus.

NECRODES SURINAMENSIS (SILPHIDAE) IN WESTERN WASHINGTON

Hatch (1957)¹ suggests that *Necrodes surinamensis* has recently been introduced "west of the Rocky Mountains where it is rapidly becoming established." In the Pacific Northwest Hatch cites records of one specimen from northeastern British Columbia (no date), one from Idaho (1949), a series from eastern Washington (1949, 1951, 1956), and three from western Oregon (1946, 1955).

In May, 1958, the writer collected two adult specimens of this species from a dead raccoon found in Dosewallips Canyon, Jefferson Co., Washington. These specimens from the Olympic Peninsula establish a new western Washington record and further corroborate Dr. Hatch's theory that *Necrodes surinamensis* has invaded the Pacific Northwest, and is becoming established there.—JENS W. KNUDSEN, Dept. of Biol., Pacific Lutheran College, Parkland, Wn.

¹Hatch, Melville H. 1957. The Beetles of the Pacific Northwest. II: Staphyliniformia, pg. 11.